

ASCA Discovery of Diffuse 6.4 keV Emission Near the Sgr C Complex: A New X-ray Reflection Nebula

Hiroshi Murakami¹, Katsuji Koyama², and Masahiro Tsujimoto¹

Department of Physics, Faculty of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan; hiro@cr.scphys.kyoto-u.ac.jp, koyama@cr.scphys.kyoto-u.ac.jp, tsujimot@cr.scphys.kyoto-u.ac.jp

and

Yoshitomo Maeda

Department of Astronomy and Astrophysics, The Pennsylvania State University, University Park, PA 16802-6305, U.S.A.; maeda@astro.psu.edu

and

Masaaki Sakano

Space Utilization Research Program, National Space Development Agency of Japan, 2-1-1, Sengen, Tsukuba, Ibaraki, 305-8505, Japan; sakano.masaaki@nasda.go.jp

ABSTRACT

We present an *ASCA* discovery of diffuse hard X-ray emission from the Sgr C complex with its peak in the vicinity of the molecular cloud core. The X-ray spectrum is characterized by a strong 6.4-keV line and large absorption. These properties suggest that Sgr C is a new *X-ray reflection nebula* which emits fluorescent and scattered X-rays via irradiation from an external X-ray source. We found no adequately bright source in the immediate Sgr C vicinity to fully account for the fluorescence. The irradiating source may be the Galactic nucleus Sgr A*, which was brighter in the past than it is now as is suggested from observations of the first *X-ray reflection nebula* Sgr B2.

¹Research Fellow of the Japan Society for the Promotion of Science (JSPS)

²CREST, Japan Science and Technology Corporation (JST), 4-1-8 Honmachi, Kawaguchi, Saitama, 332-0012, Japan

Subject headings: Galaxy: center — ISM: clouds — ISM: individual (Sgr C) — X-rays: ISM

1. Introduction

The Galactic Center (GC) region of our Galaxy is a unique concentration of stars and molecular clouds, in addition to Sgr A*, a $2.6 \times 10^6 M_\odot$ massive black hole (MBH) at the gravitational center (for a recent review, see Genzel & Eckart 1999).

The inner $\simeq 100$ pc region of the Galaxy is a site of a nuclear starburst and consequently should be populated by a variety of X-ray sources, such as X-ray binaries, supernova remnants, and young stellar clusters. *ASCA* revealed diffuse emissions of $2'$ extent centered at Sgr A* and asymmetrical structure extending $\simeq 1^\circ$ along the Galactic plane (Koyama et al. 1996; Maeda et al. 1998). The spectral characteristics were most remarkable, exhibiting many K-shell lines from highly ionized atoms, with particularly enhanced emission from He-like (6.70 keV) and H-like (6.97 keV) irons.

Early *Chandra* results confirmed the *ASCA* findings and resolved numerous X-ray structures: X-ray binaries, diffuse structures, emission from the supernova remnant Sgr A-East, emission from individual stars within the Sgr A region, and a possible X-ray counterpart of the massive black hole Sgr A* with a luminosity of only 10^{33} erg s $^{-1}$ (Baganoff et al. 2000), which is extremely low compared to typical massive black holes.

Among these many X-ray objects, the most massive molecular cloud Sgr B2 in the Galaxy is extraordinary. It exhibits a very peculiar spectrum with a strong emission line at 6.4 keV, a low energy cutoff below 4 keV and a pronounced edge-structure at 7.1 keV. The X-ray image is shifted from the core of the molecular cloud toward the GC by about $1\frac{1}{3}$ (Murakami et al. 2000a). Recently *Chandra*

confirmed these results: the extended structure of the 6.4 keV line image with a convex shape pointed towards Sgr A* and its offset morphology from the cloud core (Murakami et al. 2000b). A numerical simulation demonstrated that the offset morphology and the line and edge dominated spectrum are well reproduced by the reflection of external X-rays coming from the direction of the GC (Figure 2, Murakami et al. 2000a; also in Sunyaev et al. 1998). Based on this, they proposed Sgr B2 to be a new class of X-ray object, an *X-ray reflection nebula* (XRN). However, there is no adequately bright source in the immediate Sgr B2 vicinity to fully account for the fluorescence. They suspected that the most likely source is an X-ray outburst from the MBH at Sgr A* despite its considerable distance from Sgr B2 (Koyama et al. 1996; Murakami et al. 2000a). Their model requires an outburst of $L_X \sim 10^{39}$ erg s $^{-1}$ for at least 10 years and must have stopped at most 30 years ago to avoid detection by earlier X-ray astronomical instruments. Such an outburst from a $10^6 M_\odot$ MBH, perhaps caused by a surge in accretion rate, is consistent with the behavior of active galactic nuclei.

The XRN scenario and the putative past X-ray outburst of the Galactic nucleus Sgr A* may indicate that there should be many other XRN in the GC region. We therefore examined the archival *ASCA* data of the GC region in detail, and found another XRN candidate in the Sgr C region, located at the same projected distance as Sgr B2 from the GC, but in the opposite direction.

2. Observation

The *ASCA* observations of Sgr C were made on 1993 October 4. All four focal plane instruments, two Solid-state Imaging Spec-

trometers (SIS0, SIS1) and two Gas Imaging Spectrometers (GIS2, GIS3) were operated in parallel, providing four independent data sets. Details of the instruments, the telescopes and the detectors, are found in Tanaka, Inoue, & Holt (1994), Serlemitsos et al. (1995), Burke et al. (1991), Ohashi et al. (1996), Makishima et al. (1996), and Gotthelf (1996). Each of the GISs was operated in PH mode with the standard bit-assignment, while the SIS was operated in 4-CCD bright mode. The data were post-processed to correct for spatial gain non-linearity. Data taken at geomagnetic cutoff rigidities lower than 6 GV, at elevation angles less than 5° from the earth rim, or during the passage through the South Atlantic Anomaly were excluded. After these filterings, the net observing time was 20 ksec.

3. Analysis and Results

The key characteristic of XRN is a strong iron line at 6.4 keV (Koyama et al. 1996; Sunyaev & Churazov 1998; Murakami et al. 2000a), hence we made a narrow energy band image with a central energy of 6.4 keV and a width of twice the energy resolution (FWHM): 5.8–7.0 keV for the GIS. Figure 1 shows the GIS image overlaid with the contour map of the cold cloud density distribution contour by the radio observation of CS with the radial velocity of -120 km s^{-1} to -110 km s^{-1} (Figure 3 in Tsuboi, Handa, & Ukita 1999). The position accuracy is about $5''$ for the radio map, and $24''$ for the GIS image (Gotthelf et al. 2000). The 6.4 keV band image shows an X-ray peak (inner region of the solid circle) near the molecular cloud core CO 359.4 + 0.0 (Oka et al. 1998), which is one of the giant molecular clouds in the Sgr C complex, with a mass of $\sim 10^6 M_\odot$. We therefore suspect that the X-ray emission is associated with the

molecular cloud Sgr C, although a more detailed comparison will require more accurate positional observations both in X-rays and radio bands.

We made X-ray spectra using the X-ray photons in a circle of $2'.7$ radius around the X-ray peak (see Figure 1). Since the Sgr C region is in the large scale GC plasma (Koyama et al. 1996), the spectrum may be contaminated by the emission of He- and H-like iron lines (6.70 keV, and 6.97 keV). In order to properly subtract these highly ionized iron lines, the background region is selected as a $2'.7$ radius region with the same Galactic latitude (dotted circle in Figure 1).

Figure 2 shows the background subtracted GIS spectrum. We made a model spectrum with a power-law continuum and a Gaussian line, folding with the *ASCA* response function, and fitted it to the data. Due to the limited statistics, we fixed the power-law photon index α to be 2.0 (number of photons $N(E)dE \propto E^{-\alpha}dE$, where E is the photon energy), the same value as assumed for Sgr B2 (Koyama et al. 1996; Murakami et al. 2000a). The fit is acceptable, with the best-fit parameters shown in Table 1. The central energy of the line appears at $6.28^{+0.15}_{-0.21}$ keV, in agreement with the fluorescent line from neutral irons. The line equivalent width and absorption column density are both very large: $\sim 0.8 \text{ keV}$ and $1.3^{+0.3}_{-0.4} \times 10^{23} \text{ H cm}^{-2}$, respectively. Since the Galactic diffuse background may not be uniform, we checked the ambiguity of the background selection. We selected three other background regions with the same Galactic latitude as the source region, and fitted them with the same model. The best-fit parameters of the line central energy and the absorption are in the range of statistical errors. The absorbed flux differs a little larger

than the error, however this flux variation is not serious for the discussion in section 4.

Adding two narrow Gaussian lines at 6.4 and 6.7 keV on a power-law continuum, we found that the 6.7-keV line from highly ionized iron is very weak; the upper limit of $9 \times 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$ is about 1/4 of the best-fit value of the 6.4-keV line. We thus conclude that the emission line from the Sgr C cloud is dominated by the fluorescent line from cold irons.

We also made an SIS spectrum from the same region of GIS, and fitted it with the same model as with the GIS. The best-fit spectral parameters are shown in Table 1, and are consistent with those obtained from the GIS spectrum. Due to the limited statistics, the SIS spectrum does not further constrain the spectral parameters. We thus discuss the X-ray properties of Sgr C using the spectral parameters obtained with GIS only.

4. Discussion

4.1. Is Sgr C a New XRN?

We have found a hard X-ray enhancement near the Sgr C cloud on the 6.4 keV line map. The spectrum shows a strong fluorescent line at 6.4 keV and the large absorption in low energy band. These properties are good evidence that Sgr C is a new XRN (Sunyaev & Churazov 1998; Murakami et al. 2000a).

Murakami et al. (2000a) analyzed the spectrum of Sgr B2, the XRN, and found that the equivalent width of the 6.4 keV line is $2.9_{-0.9}^{+0.3} \text{ keV}$, and the column density of hydrogen is $8.3_{-2.0}^{+2.5} \times 10^{23} \text{ H cm}^{-2}$. Those of Sgr C are $0.8_{-0.5}^{+0.4} \text{ keV}$ and $1.3_{-0.4}^{+0.3} \times 10^{23} \text{ H cm}^{-2}$, respectively.

The absorption of Sgr C is smaller than that of Sgr B2 by an order of magnitude. This

is reasonable because the mass of the Sgr C cloud is about 1/7 of the Sgr B2 cloud (Oka et al. 1998), in spite of rather similar geometrical size. In fact, using the Sgr C cloud size of $\sim 28 \text{ pc}$ (Oka et al. 1998), we can roughly estimate the column density to be $4 \times 10^{22} \text{ H cm}^{-2}$, which is almost equal to $1.3 \times 10^{23} \text{ H cm}^{-2}$, after taking account of the interstellar absorption to the Sgr C cloud of $\sim 10^{23} \text{ H cm}^{-2}$ (Sakano et al. 1998).

The absorption column of $4 \times 10^{22} \text{ H cm}^{-2}$ is in the optically thin range near 6.4 keV and above 7.1 keV energy, hence, unlike Sgr B2, the 6.4 keV emission region should overlap the molecular core. This is in agreement with the observation, although the observational results still have significant uncertainty.

4.2. Irradiating Source of XRN

In the optically thin case, the X-ray intensity of an XRN is simply proportional to the amount of the scattering matter and the flux of the irradiating X-rays. The absorption-corrected luminosity of the 6.4 keV line from Sgr C is $4 \times 10^{33} \text{ erg s}^{-1}$ within a 2.7 -radius circular region. The required luminosity of the X-ray source irradiating Sgr C would be about $3 \times 10^{39} (d_{\text{SgrC}}/100 \text{ pc})^2 \text{ erg s}^{-1}$, where d_{SgrC} is the distance from the irradiating X-ray source to Sgr C.

The brightest nearby X-ray source is 1E 1740.7–2942, which lies only ~ 0.4 degrees from Sgr C. During the present observation, its luminosity was $\sim 3 \times 10^{36} \text{ erg s}^{-1}$ in the 2–10 keV band (Sakano et al. 1999). This luminosity, however, is two orders of magnitude less than that required to account for the fluorescent X-rays from Sgr C. Even if we accumulate the X-ray fluxes from all cataloged bright X-ray sources near the GC (Sakano 2000), we can explain only 2 % of the reflected luminosity of Sgr C.

This fact strengthens the scenario that the X-rays from Sgr C, as well as those from Sgr B2, is due to the reflection from an irradiating source which was very bright in the past but is presently dim. The X-ray fluxes of Sgr B2 and Sgr C can be consistently explained by one irradiating source, if it is located at almost the same distance from both of the XRN. This position falls near the Galactic nucleus Sgr A*, and the X-ray luminosity should be as luminous as 3×10^{39} erg s⁻¹ about 300 years ago, the light travel time from Sgr A* to Sgr B2 and Sgr C.

5. Summary

1. We have discovered diffuse emission in the 6.4 keV band from the Sgr C region based on the *ASCA* observations.

2. We found that the X-ray spectrum of Sgr C shows a strong emission line at 6.4 keV from cold iron with an equivalent width of about 0.8 keV, and a large absorption of 1.3×10^{23} H cm⁻².

3. The line dominated spectrum and the image correlated with a molecular cloud are similar to those of the prototypical X-ray reflection nebula (XRN) Sgr B2, and thus Sgr C can be regarded as a new XRN.

4. There is no adequately bright source in the immediate Sgr C vicinity to fully account for the observed fluorescence.

5. Existence of the two XRN, Sgr B2 and Sgr C, can be fully explained if we consider the AGN activity some hundreds of years ago at the GC ($L_X \sim 3 \times 10^{39}$ erg s⁻¹).

ety for the Promotion of Science. Thanks are also due to Patrick Durrell for a critical reading of the manuscript.

The authors express their thanks to all the members of the *ASCA* team. H.M and M.T. are financially supported by the Japan Soci-

REFERENCES

- Baganoff, F., et al. 2000, Astronomical Society of Japan Meeting, R31c
- Burke, B. E., Mountain, R. W., Harrison, D. C., Bautz, M. W., Doty, J. P., Ricker, G. R., & Daniels, P. J. 1991, IEEE Trans., ED-38, 1069
- Genzel, R., & Eckart, A. 1999, The Central Parsecs of the Galaxy, ASP Conference Series, 186, 3
- Gotthelf, E. 1996, The *ASCA* news (Greenbelt: NASA GSFC), 4, 31
- Gotthelf, E. V., Ueda, Y., Fujimoto, R., Kii, T., & Yamaoka, K. 2000, ApJ, 543, 417
- Koyama, K., Maeda, Y., Sonobe, T., Takeshima, T., Tanaka, Y., & Yamauchi, S. 1996, PASJ, 48, 249
- Maeda, Y., Koyama, K., Murakami, H., Sakano, M., Ebisawa, K., Takeshima, T., & Yamauchi, S. 1998, in IAU Symp. 188, The Hot Universe, eds. K. Koyama, S. Kitamoto, M. Itoh (London: Kluwer), 352
- Makishima, K., et al. 1996, PASJ, 48, 171
- Murakami, H., Koyama, K., Sakano, M., Tsujimoto, M., & Maeda, Y. 2000a, ApJ, 534, 238
- Murakami, H., Maeda, Y., Sakano, M., Koyama, K., Tsujimoto, M. 2000b, American Astronomical Society Meeting, 196, 5902
- Ohashi, T., et al. 1996, PASJ, 48, 157
- Oka, T., Hasegawa, T., Hayashi, M., Hanada, T., & Sakamoto, S. 1998, ApJ, 493, 730
- Sakano, M., Nishiuchi, M., Maeda, Y., Koyama, & Yokogawa, J. 1998, IAU Symp. 184, The Central Regions of the Galaxy and Galaxies, ed. Y. Sofue (London: Kluwer), 443
- Sakano, M., Imanishi, K., Tsujimoto, M., Koyama, K., & Maeda, Y. 1999, ApJ, 520, 316
- Sakano, M. 2000, Ph.D. thesis, Kyoto University
- Serlemitsos, P. J., et al. 1995, PASJ, 47, 105
- Sunyaev, R., & Churazov, E. 1998, MNRAS, 297, 1279
- Tanaka, Y., Inoue, H., & Holt, S. S. 1994, PASJ, 46, L37
- Tsuboi, M., Handa, T., & Ukita, N. 1999, ApJS, 120, 39

This 2-column preprint was prepared with the AAS L^AT_EX macros v4.0.




Fig. 1.— The 5.8–7.0 keV band image around the Sgr C cloud obtained with the GIS, laid over the CS line contours (Figure 3 in Tsuboi, Handa, & Ukita 1999). The solid and the dotted circle show the source and the background regions, respectively.




Fig. 2.— The GIS (GIS2 + 3) spectrum of Sgr C. The solid line shows the best-fit model shown in Table 1.

TABLE 1
FITTING RESULTS OF SGR C TO A PHENOMENOLOGICAL SPECTRAL MODEL

Model Components	Parameters	Unit	GIS	SIS
Absorption	N_{H}^{a}	$(10^{22} \text{ H cm}^{-2})$	$12.6^{+3.5}_{-3.3}$	$9.7^{+3.9}_{-2.5}$
Continuum	Photon Index		2.0 (fixed)	2.0 (fixed)
	Flux ^b (2–10 keV)	$(10^{-4} \text{ ph s}^{-1} \text{ cm}^{-2})$	$3.0^{+0.9}_{-0.6}$	$2.0^{+0.7}_{-0.5}$
Fe Line	Center Energy	(keV)	$6.28^{+0.15}_{-0.21}$	$6.38^{+0.10}_{-0.31}$
	Flux ^b	$(10^{-5} \text{ ph s}^{-1} \text{ cm}^{-2})$	$3.5^{+1.4}_{-2.2}$	$2.7^{+2.3}_{-1.5}$
	Equivalent Width	(keV)	$0.8^{+0.4}_{-0.5}$	$1.1^{+1.1}_{-0.7}$
Total Luminosity	$L_{4-10\text{keV}}$	$(10^{34} \text{ erg s}^{-1})$	5.0	3.0
Reduced χ^2 (d.o.f.)			0.80 (20)	0.63 (14)

NOTE.—The errors are at 90% confidence level.

^aThe equivalent hydrogen column density for the solar abundances.

^bThe fluxes are not corrected for absorption.